

A NOTE ON WEIGHTED HOMOGENEOUS SICIAK-ZAHARYUTA EXTREMAL FUNCTIONS

BARBARA DRINOVEC DRNOVŠEK AND RAGNAR SIGURDSSON

ABSTRACT. We prove that for any given upper semicontinuous function φ on an open subset E of $\mathbb{C}^n \setminus \{0\}$, such that the complex cone generated by E minus the origin is connected, the homogeneous Siciak-Zaharyuta function with the weight φ on E , can be represented as an envelope of a disc functional.

Introduction. Let \mathcal{L} denote the Lelong class on \mathbb{C}^n and \mathcal{L}^h the subclass of functions u which are *logarithmically homogeneous*. Let $\varphi: E \rightarrow \overline{\mathbb{R}}$ be a function on a subset E of \mathbb{C}^n taking values in the extended real line $\overline{\mathbb{R}}$. The *Siciak-Zaharyuta extremal function* $V_{E,\varphi}$ with weight φ is defined by

$$V_{E,\varphi} = \sup\{u \in \mathcal{L}; u|_E \leq \varphi\}.$$

The *homogeneous Siciak-Zaharyuta extremal function* $V_{E,\varphi}^h$ with weight φ is defined similarly with \mathcal{L}^h in the role of \mathcal{L} . In the special case when $\varphi = 0$ we only write V_E (and V_E^h) and we call this function the (*homogeneous*) *Siciak-Zaharyuta extremal function for the set E* . The function V_E (V_E^h) is also called the (*homogeneous*) *pluricomplex Green function for E with pole at infinity*.

Theorem 1. *Let $\varphi: E \rightarrow \mathbb{R} \cup \{-\infty\}$ be an upper semicontinuous function on an open subset E of $\mathbb{C}^n \setminus \{0\}$. Assume that there exists a function in \mathcal{L}^h dominated by φ on E . Then the largest logarithmically homogeneous function $\mathbb{C}E \rightarrow \mathbb{R} \cup \{-\infty\}$ dominated by φ on E is upper semicontinuous on \mathbb{C}^*E and it is of the form $\log \varrho_{E,\varphi}$, where*

$$(1) \quad \varrho_{E,\varphi}(z) = \inf\{|\lambda|e^{\varphi(z/\lambda)}; \lambda \in \mathbb{C}^*, z/\lambda \in E\}, \quad z \in \mathbb{C}^*E.$$

*If \mathbb{C}^*E is connected, then for every $z \in \mathbb{C}^n$*

$$(2) \quad V_{E,\varphi}^h(z) = \inf \left\{ \int_{\mathbb{T}} \log \varrho_{E,\varphi}(f_1, \dots, f_n) d\sigma; f \in \mathcal{O}(\overline{\mathbb{D}}, \mathbb{P}^n), f = [f_0 : \dots : f_n], \right. \\ \left. f(\mathbb{T}) \subset \mathbb{C}^*E, f_0(0) = 1, (f_1(0), \dots, f_n(0)) = z. \right\}$$

Date: February 2, 2015.

2010 Mathematics Subject Classification. Primary 32U35; Secondary 32U15, 32U05.

Key words and phrases. Siciak's homogeneous extremal function, envelope of a disc functional.

B. Drinovec Drnovšek was partially supported by grant P1-0291, Republic of Slovenia.

If $\mathbb{C}E = \mathbb{C}^n$, then for every $z \in \mathbb{C}^n$

$$(3) \quad V_{E,\varphi}^h(z) = \inf \left\{ \int_{\mathbb{T}} \log \varrho_{E,\varphi} \circ f \, d\sigma ; f \in \mathcal{O}(\overline{\mathbb{D}}, \mathbb{C}^n), f(0) = z \right\}.$$

A *disc envelope formula* is a formula where the values of a function F defined on a complex space X with values on the extended real line $\overline{\mathbb{R}}$ are given as $F(z) = \inf\{H(f); f \in \mathcal{B}(z)\}$, where H is *disc functional*, i.e., a function defined on some subset \mathcal{A} of $\mathcal{O}(\mathbb{D}, X)$, the set of *analytic discs* in X , with values on $\overline{\mathbb{R}}$, \mathcal{B} is a subclass of \mathcal{A} , and $\mathcal{B}(z)$ consists of all of $f \in \mathcal{B}$ with *center* $z = f(0)$.

The formula (2) is an example of a disc envelope formula, where \mathcal{A} consists of all closed analytic discs with value in the projective space, i.e., elements f in $\mathcal{O}(\overline{\mathbb{D}}, \mathbb{P}^n)$ which map the unit circle \mathbb{T} into \mathbb{C}^*E , $H(f)$ is the integral, and \mathcal{B} is the subset of \mathcal{A} consisting of discs with $f_0(0) = 1$. We identify a point $[1 : z] \in \mathbb{P}^n$ with the point $z \in \mathbb{C}^n$.

For general information on the Siciak-Zaharyuta extremal function see Siciak [8, 9, 10, 11, 12] and Zaharyuta [13]. The first disc envelope formula for V_E was proved by Lempert in the case when E is an open convex subset of \mathbb{C}^n with real analytic boundary. (The proof is given in Momm [5, Appendix].) Lárusson and Sigurdsson [2] proved disc envelope formulas for V_E for open connected subsets E of \mathbb{C}^n . Magnússon and Sigurdsson [4] generalized this result and obtained a disc formula for $V_{E,\varphi}$ in the case when φ is an upper semicontinuous function on an open connected subset E of \mathbb{C}^n . Drinovec Drnovšek and Forstnerič [1] proved disc envelope formulas for V_E for open subsets E of an irreducible and locally irreducible algebraic subvariety of \mathbb{C}^n . Recently, Magnússon [3] established disc envelope formulas for the global extremal function in projective space.

Acknowledgement. This paper was written while the second author was visiting University of Ljubljana in the autumn of 2012. He would like to thank the members of the Department of Mathematics for their great hospitality and for many interesting and helpful discussions.

Notation. Let \mathbb{D} denote the unit disc in \mathbb{C} , \mathbb{T} the unit circle, and σ the arc length measure on \mathbb{T} normalized to 1. An analytic disc is a holomorphic map $f: \mathbb{D} \rightarrow X$, where X is some complex space. We let $\mathcal{O}(\mathbb{D}, X)$ denote the set of all analytic discs. We say that the disc is closed if it extends as a holomorphic map to some neighbourhood of the closed unit disc $\overline{\mathbb{D}}$ with values in X and we let $\mathcal{O}(\overline{\mathbb{D}}, X)$ denote the set of all closed analytic discs in X . The point $z = f(0) \in X$ is called the center of f .

For a subset X of \mathbb{C}^n we let $\mathcal{USC}(X)$ denote the set of all upper semicontinuous functions on X , and for open subset U of \mathbb{C}^n we denote by $\mathcal{PSH}(U)$ the set of all plurisubharmonic functions on U . The Lelong class \mathcal{L} consists of all $u \in \mathcal{PSH}(\mathbb{C}^n)$ such that $u - \log^+ |\cdot|$ is bounded above and \mathcal{L}^h is the

subclass of all logarithmically homogeneous functions, i.e., functions satisfying $u(\lambda z) = u(z) + \log |\lambda|$ for $\lambda \in \mathbb{C}^*$ and $z \in \mathbb{C}^n$. Observe that every such function takes the value $-\infty$ at the origin. For every subset E of \mathbb{C}^n we set $\mathbb{C}E = \{\lambda z; \lambda \in \mathbb{C}, z \in E\}$, $\mathbb{C}^*E = \{\lambda z; \lambda \in \mathbb{C}^*, z \in E\}$ and we call $\mathbb{C}E$ the complex cone generated by E . Note that complex cones are suitable sets for the domains of definition of logarithmically homogeneous functions.

Let \mathbb{P}^n denote the n -dimensional projective space, $\pi: \mathbb{C}^{n+1} \setminus \{0\} \rightarrow \mathbb{P}^n$ the natural projection, $(z_0, \dots, z_n) \mapsto [z_0 : \dots : z_n]$, and identify \mathbb{C}^n with the subset of all $[z_0 : \dots : z_n]$ with $z_0 \neq 0$ and, in particular, the point $z \in \mathbb{C}^n$ with $[1 : z] \in \mathbb{P}^n$. The hyperplane at infinity is $H_\infty = \pi(Z_0 \setminus \{0\})$, where Z_0 is the hyperplane in \mathbb{C}^{n+1} defined by the equation $z_0 = 0$. Then $\mathbb{P}^n = \mathbb{C}^n \cup H_\infty$.

Review of a few results. Assume that $\psi: X \rightarrow \mathbb{R} \cup \{-\infty\}$ is a measurable function on a subset X of \mathbb{C}^n , such that there is $u \in \mathcal{L}$ satisfying $u|X \leq \psi$. It is an easy observation that a function $u \in \mathcal{PSH}(\mathbb{C}^n)$ is in \mathcal{L} if and only if the function

$$(z_0, \dots, z_n) \mapsto u(z_1/z_0, \dots, z_n/z_0) + \log |z_0|$$

extends as a plurisubharmonic function from $\mathbb{C}^{n+1} \setminus Z_0$ to $\mathbb{C}^{n+1} \setminus \{0\}$. Let v denote this extension. Take $f = [f_0 : \dots : f_n] \in \mathcal{O}(\overline{\mathbb{D}}, \mathbb{P}^n)$ with $f_0(0) = 1$, $(f_1(0), \dots, f_n(0)) = z$, satisfying $f(\mathbb{T}) \subset X$, and define $\tilde{f} = (f_0, \dots, f_n) \in \mathcal{O}(\overline{\mathbb{D}}, \mathbb{C}^{n+1} \setminus \{0\})$. Then the subaverage property of $v \circ \tilde{f}$ and the Riesz representation formula applied to $\log |f_0|$ give (see [4, p. 243])

$$\begin{aligned} u(z) &= u(f_1(0), \dots, f_n(0)) + \log |f_0(0)| = v \circ \tilde{f}(0) \\ &\leq \int_{\mathbb{T}} u(f_1/f_0, \dots, f_n/f_0) d\sigma + \int_{\mathbb{T}} \log |f_0| d\sigma \\ &\leq \int_{\mathbb{T}} \psi(f_1/f_0, \dots, f_n/f_0) d\sigma - \sum_{a \in f^{-1}(H_\infty)} m_{f_0}(a) \log |a|. \end{aligned}$$

For an open connected $X \subset \mathbb{C}^n$ and $\psi \in \mathcal{USC}(X)$, Magnússon and Sigurdsson [4, Theorem 2] proved that for every $z \in \mathbb{C}^n$

$$\begin{aligned} V_{X,\psi}(z) &= \inf \left\{ - \sum_{a \in f^{-1}(H_\infty)} m_{f_0}(a) \log |a| + \int_{\mathbb{T}} \psi(f_1/f_0, \dots, f_n/f_0) d\sigma; \right. \\ (4) \quad &\left. f \in \mathcal{O}(\overline{\mathbb{D}}, \mathbb{P}^n), f(\mathbb{T}) \subset X, f_0(0) = 1, (f_1(0), \dots, f_n(0)) = z \right\}. \end{aligned}$$

Our main result, Theorem 1, will follow from this formula and the following

Proposition 2. *Let $\varphi: E \rightarrow \mathbb{R} \cup \{-\infty\}$ be a function on a subset $E \subset \mathbb{C}^n \setminus \{0\}$ such that there exists $u \in \mathcal{L}^h$ satisfying $u|E \leq \varphi$. Let $\tilde{\varphi}: \mathbb{C}E \rightarrow \mathbb{R} \cup \{-\infty\}$ be the supremum of all logarithmically homogeneous functions on $\mathbb{C}E$ dominated by φ on E . Then the following hold:*

- (i) $\tilde{\varphi}$ is logarithmically homogeneous on \mathbb{C}^*E and for every $z \in \mathbb{C}^*E$
- (5)
$$\tilde{\varphi}(z) = \inf\{\varphi(\lambda z) - \log |\lambda|; \lambda \in \mathbb{C}^* \text{ and } \lambda z \in E\},$$
- (ii) $V_{E,\varphi}^h = V_{E,\tilde{\varphi}}^h = V_{\mathbb{C}^*E,\tilde{\varphi}}^h.$

If, in addition to the above, \mathbb{C}^*E is nonpluripolar and $\varphi \in \mathcal{USC}(E)$ then

- (iii) $\tilde{\varphi} \in \mathcal{USC}(\mathbb{C}^*E)$ and $V_{E,\varphi}^h = V_{\mathbb{C}^*E,\tilde{\varphi}}^h,$
- (iv) if $\mathbb{C}E = \mathbb{C}^n$, then $\tilde{\varphi} \in \mathcal{USC}(\mathbb{C}^n)$ and

$$V_{E,\varphi}^h = \sup\{u \in \mathcal{PSH}(\mathbb{C}^n); u \leq \tilde{\varphi}\}.$$

Proof. (i) It is easy to see that the supremum of any family of logarithmically homogeneous functions defined on a complex cone is a logarithmically homogeneous function provided the family is bounded from above at any point of the cone. Take $z \in \mathbb{C}^*E$ and choose $\lambda \in \mathbb{C}^*$ such that $\lambda z \in E$. For any logarithmically homogeneous function u on $\mathbb{C}E$ dominated by φ on E we have

$$(6) \quad u(z) = u(\lambda z) - \log |\lambda| \leq \varphi(\lambda z) - \log |\lambda|$$

which implies that the family is bounded from above at z . Since all logarithmically homogeneous functions take the value $-\infty$ at the origin the family is bounded from above at any point of the cone.

Let ψ denote the function on \mathbb{C}^*E whose value at z is given by the right hand side of the equation (5). For a logarithmically homogeneous function u on $\mathbb{C}E$, dominated by φ on E , we have $u(z) \leq \varphi(\lambda z) - \log |\lambda|$ for any $\lambda \in \mathbb{C}^*$ such that $\lambda z \in E$ by (6). Taking infimum over all $\lambda \in \mathbb{C}^*$ with $\lambda z \in E$ shows that $u \leq \psi$ on \mathbb{C}^*E . Hence $\tilde{\varphi} \leq \psi$ on \mathbb{C}^*E . To prove the converse inequality note that

$$(7) \quad \begin{aligned} \psi(\mu z) &= \inf\{\varphi(\lambda \mu z) - \log |\lambda|; \lambda \in \mathbb{C}^* \text{ and } \lambda \mu z \in E\} \\ &= \inf\{\varphi(\lambda \mu z) - \log |\lambda \mu|; \lambda \in \mathbb{C}^* \text{ and } \lambda \mu z \in E\} + \log |\mu| \\ &= \psi(z) + \log |\mu| \end{aligned}$$

for any $z \in \mathbb{C}^*E$ and $\mu \in \mathbb{C}^*$ thus the map ψ is logarithmically homogeneous. Since $\psi \leq \varphi$ on E we get $\psi \leq \tilde{\varphi}$.

(ii) Since $\varphi \geq \tilde{\varphi}$ on E and $E \subset \mathbb{C}^*E$ we have $V_{E,\varphi}^h \geq V_{E,\tilde{\varphi}}^h \geq V_{\mathbb{C}^*E,\tilde{\varphi}}^h$. For proving the two equalities we take $u \in \mathcal{L}^h$ with $u|_E \leq \varphi$. By (i) we obtain $u \leq \tilde{\varphi}$ on \mathbb{C}^*E which implies $V_{\mathbb{C}^*E,\tilde{\varphi}}^h \geq V_{E,\varphi}^h$.

(iii) To prove that $\tilde{\varphi}$ is upper semicontinuous take $z_0 \in \mathbb{C}^*E$ and $c > \tilde{\varphi}(z_0)$. We need to show that $c > \tilde{\varphi}(z)$ for all z in some neighbourhood U of z_0 . We choose $\lambda_0 \in \mathbb{C}^*$ such that $\lambda_0 z_0 \in E$ and such that $c > \varphi(\lambda_0 z_0) - \log |\lambda_0|$. Since $\varphi \in \mathcal{USC}(E)$ there exists an open neighbourhood U of z_0 such that $\lambda_0 z \in E$ and $c > \varphi(\lambda_0 z) - \log |\lambda_0|$ for all $z \in U$. By (i) we have $c > \tilde{\varphi}(z)$ for all $z \in U$.

Since $\mathcal{L}^h \subset \mathcal{L}$ we have $V_{\mathbb{C}^*E,\tilde{\varphi}}^h \leq V_{\mathbb{C}^*E,\tilde{\varphi}}$. For proving the opposite inequality we take $u \in \mathcal{L}$ such that $u \leq \tilde{\varphi}$ on \mathbb{C}^*E . Then $u(\lambda z) - \log |\lambda| \leq \tilde{\varphi}(\lambda z) - \log |\lambda| = \tilde{\varphi}(z)$ for all $z \in \mathbb{C}^*E$ and $\lambda \in \mathbb{C}^*$. Let v be the upper

semicontinuous regularization of the function $\sup\{u(\lambda \cdot) - \log |\lambda|; \lambda \in \mathbb{C}^*\}$ on \mathbb{C}^n . We have $u \leq v \leq \tilde{\varphi}$ on \mathbb{C}^*E and since \mathbb{C}^*E is nonpluripolar and $\tilde{\varphi}$ is locally bounded above on \mathbb{C}^*E , we have $v \in \mathcal{L}$. A similar calculation as in (7) shows that v is logarithmically homogeneous, which proves the opposite inequality.

(iv) The fact that $\tilde{\varphi}$ is upper semicontinuous at 0 easily follows from the fact that $\tilde{\varphi}$ is bounded from above on the unit sphere and that it is logarithmically homogeneous. By (iii) we get $V_{E,\varphi}^h = V_{\mathbb{C}^*E,\tilde{\varphi}}$ and it is easy to see that in the case $\mathbb{C}E = \mathbb{C}^n$ the latter equals $V_{\mathbb{C}^n,\tilde{\varphi}}$.

Let $P_{\tilde{\varphi}}$ denote the function whose value at z is given by the right hand side of the equation. Since $\mathcal{L} \subset \mathcal{PSH}(\mathbb{C}^n)$ it follows $V_{\mathbb{C}^n,\tilde{\varphi}} \leq P_{\tilde{\varphi}}$. To prove the opposite inequality, it is enough to show that $P_{\tilde{\varphi}} \in \mathcal{L}$. Since $\tilde{\varphi} \in \mathcal{USC}(\mathbb{C}^n)$ it follows that $P_{\tilde{\varphi}}$ is the largest plurisubharmonic function on \mathbb{C}^n dominated by $\tilde{\varphi}$. By upper semicontinuity the map $\tilde{\varphi}$ is bounded from above on the unit sphere in \mathbb{C}^n by some constant $M \in \mathbb{R}$. Since $\tilde{\varphi}$ is logarithmically homogeneous we get

$$P_{\tilde{\varphi}}(\lambda z) \leq \tilde{\varphi}(\lambda z) \leq \log |\lambda| + M = \log |\lambda z| + M$$

for any $z \in \mathbb{C}^n$, $|z| = 1$, and $\lambda \in \mathbb{C}^*$. It follows that $P_{\tilde{\varphi}} \in \mathcal{L}$. \square

Proof of Theorem 1. By Proposition 2 the largest logarithmically homogeneous function $\tilde{\varphi}: \mathbb{C}E \rightarrow \mathbb{R} \cup \{-\infty\}$ dominated by φ on E is upper semicontinuous on \mathbb{C}^*E and $\varrho_{E,\varphi} = e^{\tilde{\varphi}(z)} = \inf\{|\lambda|e^{\varphi(z/\lambda)}; \lambda \in \mathbb{C}^*, z/\lambda \in E\}$ which proves (1).

If we take $X = \mathbb{C}^*E$ and $\psi = \tilde{\varphi}$ in (4), then logarithmic homogeneity of $\tilde{\varphi}$ on \mathbb{C}^*E implies that

$$\int_{\mathbb{T}} \tilde{\varphi}(f_1/f_0, \dots, f_n/f_0) d\sigma = \int_{\mathbb{T}} \tilde{\varphi}(f_1, \dots, f_n) d\sigma - \int_{\mathbb{T}} \log |f_0| d\sigma.$$

If $f_0(0) = 1$, then the Riesz representation formula gives

$$\sum_{a \in f^{-1}(H_\infty)} m_{f_0}(a) \log |a| + \int_{\mathbb{T}} \log |f_0| d\sigma = 0,$$

which implies that the right hand side of (4) reduces to

$$V_{\mathbb{C}^*E,\tilde{\varphi}}(z) = \inf \left\{ \int_{\mathbb{T}} \tilde{\varphi}(f_1, \dots, f_n) d\sigma; f \in \mathcal{O}(\overline{\mathbb{D}}, \mathbb{P}^n), \right. \\ \left. f(\mathbb{T}) \subset \mathbb{C}^*E, f_0(0) = 1, (f_1(0), \dots, f_n(0)) = z \right\},$$

thus (2) follows from Proposition 2 (iii).

If $\mathbb{C}E = \mathbb{C}^n$ then Proposition 2 (iv) and Poletsky theorem [6, 7] imply

$$V_{E,\varphi}^h = \sup\{u \in \mathcal{PSH}(\mathbb{C}^n); u \leq \tilde{\varphi}\} \\ = \inf \left\{ \int_{\mathbb{T}} \log \varrho_{E,\varphi} \circ f d\sigma; f \in \mathcal{O}(\overline{\mathbb{D}}, \mathbb{C}^n), f(0) = z \right\}$$

which proves (3). \square

Observation. In the special case $\varphi = 0$ we write ϱ_E for $\varrho_{E,\varphi}$. The function ϱ_E is absolutely homogeneous of degree 1, i.e., $\varrho_E(z\zeta) = |z|\varrho_E(\zeta)$. Thus, if E is a balanced domain, i.e., $\overline{\mathbb{D}}E = E$, then ϱ_E is its Minkowski function.

REFERENCES

- [1] Barbara Drinovec Drnovšek and Franc Forstnerič. Disc functionals and Siciak-Zaharyuta extremal functions on singular varieties. *Ann. Polon. Math.*, 106:171–191, 2012.
- [2] Finnur Lárusson and Ragnar Sigurdsson. The Siciak-Zahariuta extremal function as the envelope of disc functionals. *Ann. Polon. Math.*, 86(2):177–192, 2005.
- [3] Benedikt Steinar Magnússon. Analytic discs, global extremal functions and projective hulls in projective space. *preprint, arXiv:1304.1704*, 2013.
- [4] Benedikt Steinar Magnússon and Ragnar Sigurdsson. Disc formulas for the weighted Siciak-Zahariuta extremal function. *Ann. Polon. Math.*, 91(2-3):241–247, 2007.
- [5] Siegfried Momm. An extremal plurisubharmonic function associated to a convex pluri-complex Green function with pole at infinity. *J. Reine Angew. Math.*, 471:139–163, 1996.
- [6] Evgeny A. Poletsky. Plurisubharmonic functions as solutions of variational problems. In *Several complex variables and complex geometry, Part 1 (Santa Cruz, CA, 1989)*, volume 52 of *Proc. Sympos. Pure Math.*, pages 163–171. Amer. Math. Soc., Providence, RI, 1991.
- [7] Evgeny A. Poletsky. Holomorphic currents. *Indiana Univ. Math. J.*, 42(1):85–144, 1993.
- [8] Józef Siciak. On an extremal function and domains of convergence of series of homogeneous polynomials. *Ann. Polon. Math.*, 10:297–307, 1961.
- [9] Józef Siciak. On some extremal functions and their applications in the theory of analytic functions of several complex variables. *Trans. Amer. Math. Soc.*, 105:322–357, 1962.
- [10] Józef Siciak. Extremal plurisubharmonic functions in \mathbb{C}^n . *Ann. Polon. Math.*, 39:175–211, 1981.
- [11] Józef Siciak. Extremal plurisubharmonic functions and capacities in \mathbb{C}^n . *Sophia Kokyuroku in Mathematics*, 14, 1982.
- [12] Józef Siciak. Sets in \mathbb{C}^N with vanishing global extremal function and polynomial approximation. *Ann. Fac. Sci. Toulouse Math. (6)*, 20(Fascicule Special):189–209, 2011.
- [13] V. P. Zaharjuta. Extremal plurisubharmonic functions, orthogonal polynomials, and the Bernšteĭn-Walsh theorem for functions of several complex variables. *Ann. Polon. Math.*, 33:137–148, 1976/77.

FACULTY OF MATHEMATICS AND PHYSICS, UNIVERSITY OF LJUBLJANA, INSTITUTE OF MATHEMATICS, PHYSICS AND MECHANICS, JADRANSKA 19, 1000 LJUBLJANA, SLOVENIA
E-mail address: `barbara.drinovec@fmf.uni-lj.si`

DEPARTMENT OF MATHEMATICS, SCHOOL OF ENGINEERING AND NATURAL SCIENCES, UNIVERSITY OF ICELAND, IS-107 REYKJAVÍK, ICELAND
E-mail address: `ragnar@hi.is`